

THE CHARACTER SHEAVES ON THE GROUP COMPACTIFICATION

XUHUA HE

ABSTRACT. We give a definition of character sheaves on the group compactification which is equivalent to Lusztig's definition in [Moscow Math. J. 4 (2004) 869-896]. We also prove some properties of the character sheaves on the group compactification.

INTRODUCTION

Let G be a connected reductive algebraic group over an algebraically closed field. In [L3], Lusztig introduced a class of $G \times G$ -varieties. We denote the varieties by $\tilde{Z}_{J,y,\delta}$. The precise definition can be found in 1.3. The group G acts diagonally on $\tilde{Z}_{J,y,\delta}$. Lusztig introduced a partition of $\tilde{Z}_{J,y,\delta}$ into finitely many G -stable pieces. The G -orbits on each piece are in one-to-one correspondence with the conjugacy classes of a certain (smaller) reductive group.

To each character sheaf on the (smaller) reductive group, we associate a G -equivariant simple perverse sheaf on the G -stable piece. We call it a character sheaf on the G -stable piece. Its perverse extension to $\tilde{Z}_{J,y,\delta}$ is a G -equivariant simple perverse sheaf on $\tilde{Z}_{J,y,\delta}$. The simple perverse sheaves obtained in this way are called the “parabolic character sheaves”. (This is a generalization of the character sheaves on the group.)

The definition above is one of the two equivalent definitions in [L3]. The other one imitates Lusztig's definition of character sheaves of the group. Roughly speaking, the second definition help us to understand the perverse extensions of character sheaves on the G -stable pieces to $\tilde{Z}_{J,y,\delta}$. A consequence of the coincidence of the two definition is that the parabolic character sheaves have the following property: any composition factor of the perverse cohomology of the restriction of a parabolic character sheaf to a G -stable piece is a character sheaf on that piece.

The varieties $\tilde{Z}_{J,y,\delta}$ include more or less as a special case the strata of the group compactification $\overline{G^1}$ of G , here G is adjoint. Therefore, we obtain a partition of $\overline{G^1}$ into finitely many G -stable pieces. Lusztig

2000 *Mathematics Subject Classification.* 20G99.

The author thanks the National Science Foundation for its support through grant DMS-0243345 (principal investigator: George Lusztig).

defined the character sheaves on $\overline{G^1}$ to be the perverse extensions of the character sheaves on the G -stable pieces of $\overline{G^1}$. One may expect that the character sheaves on the group compactification have the property analogous to the property of parabolic character sheaves that we mentioned above.

To achieve this goal, we need to understand the perverse extension to $\overline{G^1}$. The main propose of this paper is to introduce an equivalent definition that helps us to understand it.

Before introducing the definition, we first recall the definition of character sheaves of the group G . Denote by B the Borel subgroup of G . The group G is decomposed into finitely many $B \times B$ -orbits. Starting with certain local systems on a $B \times B$ -orbits, we obtain some simple perverse sheaves on G . (For more details, see [MS1, no. 5].) Although this is not Lusztig's original definition, one can see that they are equivalent directly from the definition.

Now let us come back to the group compactification $\overline{G^1}$. There are finitely many $B \times B$ -orbits on $\overline{G^1}$. The closure relation of the $B \times B$ -orbits and the local systems on the orbits were studied by Springer in [S1]. These local systems play the same role in our definition of character sheaves on $\overline{G^1}$ as the local systems on the $B \times B$ -orbits of G in the definition of character sheaves on G . Then we show that the simple perverse sheaves on $\overline{G^1}$ obtained in this way coincide with those obtained from Lusztig's definition.

Now let us discuss about the ideas of the proof of the coincidence of our definition and Lusztig's definition. We denote by $\mathcal{D}(\overline{G^1})$ the derived category of constructible sheaves on $\overline{G^1}$. To each element in $\mathcal{D}(\overline{G^1})$, we associate its support, which is a closed subvariety. The counterpart (on the "level of varieties") of the equivalence of these definitions (on the "level of sheaves") is the relation between the (closure of) G -stable pieces and the (closure of) $B \times B$ -orbits. The relation was discussed in [H1] and [H2]. In [H1, 2.7], we showed that each G -stable piece is the minimal G -stable subvariety that contains a particular $B \times B$ -orbit. In [H2, 4.3], we gave an inductive way to determine in which G -stable piece an element of a $B \times B$ -orbit is contained. In this paper, we "lift" these results from the "level of varieties" to the "level of sheaves".

The content of this paper is arranged as follows. In section 1, we recall the definitions of $\tilde{Z}_{J,y,\delta}$ and the G -stable pieces on it. We also discuss some properties of the G -stable pieces. In section 2, we recall the definition of the character sheaves on the group and discuss some properties of them. In section 3, we prove our key lemma. In section 4, we first introduce a new definition of the character sheaves of group compactification and prove that the character sheaves on the group compactification have the "nice" property according to our definition. As a consequence of the "nice" property, we show that our definition is equivalent to Lusztig's. We also obtain a property about the central

characters. Our approach can also be generalized to parabolic character sheaves. In section 5, we discuss some results on the parabolic character sheaves. We also obtain a new proof of the coincidence of Lusztig's two definitions of parabolic character sheaves.

We thank Lusztig for many useful discussions on character sheaves. We also thank Springer for some advice and comments on an earlier version of this paper.

1. THE G -STABLE PIECES

1.1. Let G be a connected, reductive algebraic group over an algebraically closed field $\mathbf{k}$. Let B be a Borel subgroup of G , $T \subset B$ be a maximal torus and B^- be the opposite Borel subgroup. Let W be the corresponding Weyl group and $(s_i)_{i \in I}$ the set of simple reflections. For $w \in W$, we denote by $\text{supp}(w) \subset I$ the set of simple roots whose associated simple reflections occur in some (or equivalently, any) reduced decomposition of w . We also choose a representative $\dot{w}$ of w in G .

For $J \subset I$, let W_J be the subgroup of W generated by $\{s_j \mid j \in J\}$ and W^J (resp. ${}^JW$) be the set of minimal length coset representatives of W/W_J (resp. $W_J \backslash W$). Let w_0^J be the unique element of maximal length in W_J . (We simply write w_0 for w_0^I .) For $J, K \subset I$, we write ${}^JW^K$ for ${}^JW \cap W^K$.

For $J \subset I$, let $P_J \supset B$ be the standard parabolic subgroup defined by J and $P_J^- \supset B^-$ be the opposite of P_J . Let $L_J = P_J \cap P_J^-$. For any $J \subset I$, let $\mathcal{P}_J$ be the set of parabolic subgroups conjugate to P_J . We simply write $\mathcal{B}$ for $\mathcal{P}_\emptyset$. For $J, K \subset I$, $w \in {}^JW^K$ and $P \in \mathcal{P}_J$, $Q \in \mathcal{P}_K$, we write $\text{pos}(P, Q) = w$ if $gPg^{-1} = P_J$, $gQg^{-1} = \dot{w}P_K\dot{w}^{-1}$ for some $g \in G$.

For any parabolic subgroup P , we denote by U_P its unipotent radical. We simply write U for U_B and U^- for U_{B^-} . We denote by H_P the inverse image of the connected center of P/U_P under $P \rightarrow P/U_P$.

For any closed subgroup H of G , we denote by H_{diag} the image of the diagonal embedding of H in $G \times G$.

1.2. Let $\hat{G}$ be a possibly disconnected reductive algebraic group over $\mathbf{k}$ with identity component G . For $g \in \hat{G}$ and $H \subset G$, we write ${}^gH$ for gHg^{-1} . Let G^1 be a connected component of $\hat{G}$. There exists an isomorphism $\delta : W \rightarrow W$ such that $\delta(I) = I$ and ${}^gP \in \mathcal{P}_{\delta(J)}$ for $g \in G^1$ and $P \in \mathcal{P}_J$. We choose $g_0 \in G^1$ in the same way as [H2, 1.2]. In particular, if $G^1 = G$, then $g_0 = 1$.

1.3. Let $J, J' \subset I$ and $y \in {}^{J'}W^{\delta(J)}$ such that $y\delta(J) = J'$. For $P \in \mathcal{P}_J$ and $Q \in \mathcal{P}_{J'}$, set $A_y(P, Q) = \{g \in G^1 \mid \text{pos}(Q, {}^gP) = y\}$. Set

$$\tilde{Z}_{J,y,\delta} = \{(P, Q, \gamma) \mid P \in \mathcal{P}_J, Q \in \mathcal{P}_{J'}, \gamma \in U_{P'} \backslash A_y(P, Q)/U_P\}.$$

Define the $G \times G$ action on $\tilde{Z}_{J,y,\delta}$ by $(g_1, g_2) \cdot (P, Q, \gamma) = ({}^{g_2}P, {}^{g_1}Q, g_1\gamma g_2^{-1})$. It is easy to see that $G \times G$ acts transitively on $\tilde{Z}_{J,y,\delta}$.

Set

$$\tilde{h}_{J,y,\delta} = (P_J, {}^{\dot{y}^{-1}}P_{J'}, U_{{}^{\dot{y}^{-1}}P_{J'}} g_0 U_{P_J}) \in \tilde{Z}_{J,y,\delta}.$$

For $w, v \in W$, set

$$[\widetilde{J, w, v}]_{y,\delta} = (B \times B)(\dot{w}, \dot{v}) \cdot \tilde{h}_{J,y,\delta}.$$

It is easy to see that $[\widetilde{J, wu, v\delta^{-1}(u)}]_{y,\delta} = [\widetilde{J, w, v}]_{y,\delta}$ for $u \in W_{\delta(J)}$ and

$$\tilde{Z}_{J,y,\delta} = \bigsqcup_{w \in W^{\delta(J)}, v \in W} [\widetilde{J, w, v}]_{y,\delta}.$$

For $w \in W^{\delta(J)}$, set

$$\tilde{Z}_{J,y,\delta}^w = G_{diag} \cdot [\widetilde{J, w, 1}]_{y,\delta}.$$

By [H2, 1.3 & 1.7], we have that

$$\tilde{Z}_{J,y,\delta} = \bigsqcup_{w \in W^{\delta(J)}} \tilde{Z}_{J,y,\delta}^w.$$

The subvarieties $\tilde{Z}_{J,y,\delta}^w$ are called the G -stable pieces of $\tilde{Z}_{J,y,\delta}$.

1.4. For $w \in W^{\delta(J)}$, set

$$I(J, w, \delta) = \max\{K \subset J \mid w\delta(K) = K\}.$$

In the rest of this subsection, we fix $w \in W^{\delta(J)}$ and write K for $I(J, w, \delta)$. By [H2, 1.10], we have the following results.

(1) The map $G \times ((P_K)_{diag}(L_K \dot{w}, 1) \cdot \tilde{h}_{J,y,\delta}) \rightarrow \tilde{Z}_{J,y,\delta}^w$ defined by $(g, z) \mapsto (g, g) \cdot z$ induces an isomorphism

$$G \times_{P_K} ((P_K)_{diag}(L_K \dot{w}, 1) \cdot \tilde{h}_{J,y,\delta}) \cong \tilde{Z}_{J,y,\delta}^w,$$

where P_K acts on G on the right and acts on $(P_K)_{diag}(L_K \dot{w}, 1) \cdot \tilde{h}_{J,y,\delta}$ diagonally.

(2) The map $P_K \times L_K \dot{w} g_0 \rightarrow (P_K)_{diag}(L_K \dot{w}, 1) \cdot \tilde{h}_{J,y,\delta}$ defined by $(p, l) \mapsto (plg_0^{-1}, p) \cdot \tilde{h}_{J,y,\delta}$ for $p \in P_K$ and $l \in L_K \dot{w} g_0$ induces an affine space bundle map

$$P_K \times_{L_K} L_K \dot{w} g_0 \rightarrow (P_K)_{diag}(L_K \dot{w}, 1) \cdot \tilde{h}_{J,y,\delta},$$

where L_K acts on G on the right and acts on $L_K \dot{w} g_0$ by conjugation.

Therefore, we obtain an affine space bundle map

$$\tilde{\pi}_{J,y,\delta}^w : G \times_{L_K} L_K \dot{w} g_0 \rightarrow \tilde{Z}_{J,y,\delta}^w$$

which sends (g, l) to $(glg_0^{-1}, g) \cdot \tilde{h}_{J,y,\delta}$ for $g \in G$ and $l \in L_K \dot{w} g_0$.

Let $\vartheta : \tilde{Z}_{J,y,\delta}^w \rightarrow \tilde{Z}_{K,w,\delta}^w$ be the morphism defined by

$$\vartheta((glg_0^{-1}, g) \cdot \tilde{h}_{J,y,\delta}) = (glg_0^{-1}, g) \cdot \tilde{h}_{K,w,\delta}$$

for $g \in G$, $p \in P_K$ and $l \in L_K \dot{w} g_0$. By [H2, 1.6 & 1.10], ϑ is well defined. Moreover, we have the following commuting diagram

$$\begin{array}{ccc}
G \times_{L_K} L_K \dot{w} g_0 & \xrightarrow{\tilde{\pi}_{J,y,\delta}^w} & \tilde{Z}_{J,y,\delta}^w \\
\parallel & & \downarrow \vartheta \\
G \times_{L_K} L_K \dot{w} g_0 & \xrightarrow{\tilde{\pi}_{K,w,\delta}^w} & \tilde{Z}_{K,w,\delta}^w.
\end{array}$$

Note that $w\delta(K) = K$. It is easy to see that $L_K \dot{w} g_0$ is a connected component of $N_{\hat{G}}(L_K)$. However, in general $L_K \dot{w} g_0$ is not the identity component of $N_{\hat{G}}(L_K)$ even if $\hat{G} = G = G^1$. We will see in 4.3 that the character sheaves on a (possibly) disconnected group are involved when studying character sheaves on the group compactification even if $\hat{G} = G = G^1$.

1.5. Let $J, J' \subset I$ and $y \in {}^{J'}W^{\delta(J)}$ such that $y\delta(J) = J'$. Set

$$Z_{J,y,\delta} = \{(P, Q, \gamma) \mid P \in \mathcal{P}_J, Q \in \mathcal{P}_{J'}, \gamma \in H_{P'} \backslash A_y(P, Q) / H_P\}$$

with the $G \times G$ action defined in the same way as $\tilde{Z}_{J,y,\delta}$.

As in [L3, 11.19], we may identify groups H_P/U_P (with $P \in \mathcal{P}_J$) with a single torus Δ_J independent of the choice of P . Now Δ_J acts freely on $\tilde{Z}_{J,y,\delta}$ by $t : (P, Q, \gamma) \mapsto (P, Q, \gamma z)$ where $z \in H_P$ represents $t \in D_J$. Then we may identify $Z_{J,y,\delta}$ with $\Delta_J \backslash \tilde{Z}_{J,y,\delta}$ as $G \times G$ -varieties.

Set $h_{J,y,\delta} = (P_J, {}^{y^{-1}}P_{J'}, H_{{}^{y^{-1}}P_{J'}} g_0 H_{P_J}) \in Z_{J,y,\delta}$.

For $w, v \in W$, set $[J, w, v]_{y,\delta} = (B \times B)(\dot{w}, \dot{v}) \cdot h_{J,y,\delta}$. For $w \in W^{\delta(J)}$, set $Z_{J,y,\delta}^w = G_{diag} \cdot [J, w, 1]_{y,\delta}$. Then

$$Z_{J,y,\delta} = \bigsqcup_{w \in W^{\delta(J)}, v \in W} [J, w, v]_{y,\delta} = \bigsqcup_{w \in W^{\delta(J)}} Z_{J,y,\delta}^w.$$

The subvarieties $Z_{J,y,\delta}^w$ are called the G -stable pieces of $Z_{J,y,\delta}$.

Now fix $w \in W^{\delta(J)}$. Set $K = I(J, w, \delta)$. Then as a consequence of 1.4, we obtain the following commuting diagram

$$\begin{array}{ccc}
G \times_{L_K} L_K \dot{w} g_0 / Z^0(L_J) & \longrightarrow & Z_{J,y,\delta}^w \\
\parallel & & \downarrow \vartheta \\
G \times_{L_K} L_K \dot{w} g_0 / Z^0(L_J) & \longrightarrow & Z_{K,w,\delta}^w,
\end{array}$$

where $Z^0(L_J)$ is the connected center of L_J and ϑ is defined in the similar way as in 1.4.

1.6. In this subsection, we assume that G is adjoint. The compactification $\overline{G^1}$ of G^1 is the $G \times G$ variety which is isomorphic to the wonderful compactification of G as a variety and where the $G \times G$ action is twisted by $G \times G \rightarrow G \times G$, $(g, g') \mapsto (g, g_0 g' g_0^{-1})$. By [H2, 2.1],

we have that

$$\begin{aligned}\overline{G^1} &= \bigsqcup_{J \subset I} Z_{J, w_0 w_0^{\delta(J)}, \delta} = \bigsqcup_{J \subset I} \bigsqcup_{w \in W^{\delta(J)}, v \in W} [J, w, v]_{w_0 w_0^{\delta(J)}, \delta} \\ &= \bigsqcup_{J \subset I} \bigsqcup_{w \in W^{\delta(J)}} Z_{J, w_0 w_0^{\delta(J)}, \delta}^w.\end{aligned}$$

We write $h_{J, w_0 w_0^{\delta(J)}, \delta}$ as h_J , $[J, w, v]_{w_0 w_0^{\delta(J)}, \delta}$ as $[J, w, v]$ and $Z_{J, w_0 w_0^{\delta(J)}, \delta}^w$ as Z_J^w . We call Z_J^w the G -stable pieces of $\overline{G^1}$. We denote by π_J^w the morphism $G \times_{L_{I(J, w, \delta)}} L_{I(J, w, \delta)} \dot{w} g_0 / Z^0(L_J) \rightarrow Z_J^w$ in 1.5.

2. THE CHARACTER SHEAVES ON G^1

2.1. We follow the notation of [BBD]. Let X be an algebraic variety over $\mathbf{k}$ and l be a fixed prime number invertible in $\mathbf{k}$. We write $\mathcal{D}(X)$ instead of $\mathcal{D}_c^b(X, \bar{\mathbb{Q}}_l)$. If $K \in \mathcal{D}(X)$ and A is a simple perverse sheaf on X , we write $A \dashv K$ if A is a composition factor of ${}^p H^i(K)$ for some $i \in \mathbb{Z}$. For $A, B \in \mathcal{D}(X)$, we write $A = B[\cdot]$ if $A = B[m]$ for some $m \in \mathbb{Z}$.

Let $K, K_1, K_2, \dots, K_n \in \mathcal{D}(X)$. We say that $K \in \langle K_1, K_2, \dots, K_n \rangle$ if there exist $m \geq n$ and $K_{n+1}, K_{n+2}, \dots, K_m \in \mathcal{D}(X)$ such that $K_m = K$ and for each $n+1 \leq i \leq m$, there exists $1 \leq j, k < i$ and $n_j, n_k \in \mathbb{Z}$, such that $(K_j[n_j], K_i, K_k[n_k])$ is a distinguished triangle in $\mathcal{D}(X)$. In this case, if $H_c^i(X, K_i) = 0$ for all i , then $H_c^i(X, K) = 0$. Moreover, if $A \dashv K$, then $A \dashv K_i$ for some $1 \leq i \leq n$.

If $f : X \rightarrow Y$ is a morphism of algebraic varieties, we have functors $f_!$ and f^* between the derived categories $\mathcal{D}(X)$ and $\mathcal{D}(Y)$. Therefore

(1) if $A, A_1, A_2, \dots, A_n \in \mathcal{D}(X)$ with $A \in \langle A_1, A_2, \dots, A_n \rangle$, then $f_! A \in \langle f_! A_1, f_! A_2, \dots, f_! A_n \rangle$;

(2) if $B, B_1, B_2, \dots, B_n \in \mathcal{D}(Y)$ with $B \in \langle B_1, B_2, \dots, B_n \rangle$, then $f^* B \in \langle f^* B_1, f^* B_2, \dots, f^* B_n \rangle$.

Let H be a connected algebraic group and X, Y be varieties with a free H -action on $X \times Y$. Denote by $X \times^H Y$ the quotient space. For $K_1 \in \mathcal{D}(X)$ and $K_2 \in \mathcal{D}(Y)$ such that $K_1 \boxtimes K_2$ is H -equivariant, we denote by $K_1 \odot K_2$ be the element in $\mathcal{D}(X \times^H Y)$ whose inverse image under $X \times Y \rightarrow X \times^H Y$ is $K_1 \boxtimes K_2$.

2.2. Let p be the characteristic of k and $\mathbb{Z}_{(p)}$ be the ring of rational numbers with denominator prime to p (In particular, $\mathbb{Z}_{(p)} = \mathbb{Q}$ if $p = 0$). Set $\hat{X} = \mathbb{Z}_{(p)} \otimes_{\mathbb{Z}} X / 1 \otimes_{\mathbb{Z}} X$, where X is the character group of T . If necessary, we write $\hat{X}(T)$.

Let $\mathcal{K}(T)$ be the set of isomorphism classes of Kummer local systems on T , i. e., the set of isomorphism classes of $\bar{\mathbb{Q}}_l$ -local systems $\mathcal{L}$ of rank one on T , such that $\mathcal{L}^{\otimes m} \cong \bar{\mathbb{Q}}_l$ for some integer $m \geq 1$ invertible in $\mathbf{k}$. By [MS1, 2.1], we may identify $\mathcal{K}(T)$ with $\hat{X}$. For $\xi \in \hat{X}$, we denote by $\mathcal{L}_\xi$ the corresponding Kummer local system.

For $J \subset I$, set $T_J = T/Z^0(L_J)$. We identify $\hat{X}(T_J)$ with a subgroup of $\hat{X}$, see [S1, section 1].

In the rest of this section, we recall the definition and some properties of the character sheaves on G^1 . We follow the approach in [MS1]. (In [loc. cit.] Mars and Springer dealt with the case where $G^1 = G$. The general case can be treated in a similar way.)

2.3. By the Bruhat decomposition, we have that $G^1 = \bigsqcup_{w \in W} B\dot{w}g_0B$. For $\xi \in \hat{X}$, let $\mathcal{L}_{\xi,w,\delta}$ be the inverse image of $\mathcal{L}_\xi$ under $B\dot{w}g_0B \rightarrow T$, $u\dot{w}g_0tu' \mapsto t$ for $u, u' \in U$ and $t \in T$. Then $\mathcal{L}_{\xi,w,\delta}$ is a tame local system on $B\dot{w}g_0B$. We denote by $A_{\xi,w,\delta}$ the perverse extension of $\mathcal{L}_{\xi,w,\delta}$ to G^1 . If $w\delta(\xi) = \xi$, then $\mathcal{L}_{\xi,w,\delta}$ and $A_{\xi,w,\delta}$ are equivariant for the conjugation action of B .

Define the B -action on $G \times G^1$ by $b(g, h) = (gb^{-1}, bhb^{-1})$. Let $G \times_B G^1$ be the quotient space. The map $G \times G^1 \rightarrow G^1$, $(g, h) \mapsto ghg^{-1}$ induces a proper morphism $\gamma : G \times_B G^1 \rightarrow G^1$. By standard arguments, we have the following result.

Proposition 2.4. *Let $\xi \in \hat{X}$ and A be a simple perverse sheaf on G^1 . The following conditions on A are equivalent:*

- (i) $A \dashv (\gamma)_!(\bar{\mathbb{Q}}_l \odot A_{\xi,w,\delta})$ for some $w \in W$ with $w\delta(\xi) = \xi$.
- (ii) $A \dashv (\gamma)_!(\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi,w,\delta})$ for some $w \in W$ with $w\delta(\xi) = \xi$.

2.5. Let $\mathcal{C}_\xi(G^1)$ be the set of (isomorphism classes) of simple perverse sheaves on G^1 which satisfies the equivalent conditions (i)-(ii) with respect to ξ . The simple perverse sheaves on G^1 which belong to $\mathcal{C}_\xi(G^1)$ for some $\xi \in \hat{X}$ are called character sheaves on G^1 ; they (or their isomorphism classes) form a set $\mathcal{C}(G^1)$.

Proposition 2.6 ([L1, 11.2(c)]). *Let $\xi, \eta \in \hat{X}$ with $\eta \notin W\xi$. Then*

$$\mathcal{C}_\xi(G^1) \cap \mathcal{C}_\eta(G^1) = \emptyset.$$

3. THE KEY LEMMA

In this section and the next section, we assume that G is adjoint.

3.1. Let $J \subset I$ and $w, v \in W$. Let $p_{J,w,v} : B \times B \rightarrow [J, w, v]$ be the morphism defined by $(b_1, b_2) \mapsto (b_1\dot{w}, b_2\dot{v}) \cdot h_J$ for $b_1, b_2 \in B$. By 1.3, $p_{J,wu,v\delta^{-1}(u)} = p_{J,w,v}$ for $u \in W_{\delta(J)}$. The morphism $B \times B \rightarrow T_J$, $(t_1u_1, t_2u_2) \mapsto (\dot{w}g_0)^{-1}t_1(\dot{w}g_0)\dot{v}^{-1}t_2^{-1}\dot{v}Z^0(L_J)$ factors through a morphism $pr_{J,w,v} : [J, w, v] \rightarrow T_J$.

For $\xi \in \hat{X}(T_J) \subset \hat{X}$, $pr_{J,w,v}^* \mathcal{L}_\xi = \mathcal{L}_{\xi,J,w,v}$ is a (tame) local system on $[J, w, v]$. By [MS1, 4.1.2] and [MS2, 2.2.3], $\mathcal{L}_{\xi,J,w,v}$ has weight $(w\delta(\xi), -v\xi)$ for the $B \times B$ -action and the rank one (tame) local system on $[J, w, v]$ which have a weight for the $B \times B$ -action are of the form $\mathcal{L}_{\xi,J,w,v}$ for some $\xi \in \hat{X}(T_J)$. Moreover, $\mathcal{L}_{\xi,J,wu,v\delta^{-1}(u)} = \mathcal{L}_{\delta^{-1}(u)\xi,J,w,v}$ for

$u \in W_{\delta(J)}$. If $w\delta(\xi) = v\xi$, then $\mathcal{L}_{\xi,J,w,v}$ is equivariant for the diagonal action of B .

We denote by $A_{\xi,J,w,v}$ the perverse extension of $\mathcal{L}_{\xi,J,w,v}$ to $\overline{G^1}$. By [MS2, 2.2], for $i \in \mathbb{Z}$, $\mathcal{H}^i A_{\xi,J,w,v} |_{[J',w',v']}$ is a direct sum of local system $\mathcal{L}_{\eta,J',w',v'}$, where $\eta \in \hat{X}(T_{J'})$ and $w\delta(\xi) = w'\delta(\eta)$, $v\xi = v'\eta$. It is zero unless $[J',w',v']$ is contained in the closure of $[J,w,v]$.

In other words, we have the following result.

Lemma 3.2. *Let $J \subset I$, $w, v \in W$ and $\xi \in \hat{X}(T_J)$. Set*

$$\mathcal{I}_{J,w,v,\xi} = \left\{ (J', w', v', \eta) \mid J' \subset J, \eta \in \hat{X}(T_{J'}), w, v \in W \text{ such that} \right. \\ \left. w\delta(\xi) = w'\delta(\eta), v\xi = v'\eta \text{ and } [J', w', v'] \right. \\ \left. \text{is contained in the closure of } [J, w, v] \right\}.$$

Then we have that

$$< A_{\eta,J',w',v'} >_{(J',w',v',\eta) \in \mathcal{I}_{J,w,v,\xi}} = < (i_{J',w',v'})^* \mathcal{L}_{\eta,J',w',v'} >_{(J',w',v',\eta) \in \mathcal{I}_{J,w,v,\xi}},$$

where $i_{J',w',v'} : [J', w', v'] \hookrightarrow \overline{G^1}$ is the inclusion.

3.3. For $J \subset I$, set $B_J = (B \cap L_J)/Z^0(L_J)$. For $x \in W$, set $G_{J,x} = BxB/Z^0(L_J)$ and $G'_{J,x} = B\dot{x}U_{P_{\delta(J)}^-}(B \cap L_{\delta(J)})/Z^0(L_{\delta(J)})$.

Define $\phi_{J,x} : G_{J,x} \rightarrow T_J$ by $\phi_{J,x}(u\dot{x}tu') = t$ for $t \in T_J$ and $u, u' \in U$. Define $\phi'_{J,x} : G'_{J,x} \rightarrow T_{\delta(J)}$ by $\phi'_{J,x}(u\dot{x}tu') = t$ for $t \in T_{\delta(J)}$ and $u \in U$ and $u' \in U_{P_{\delta(J)}^-}(U \cap L_{\delta(J)})$.

Let $\xi \in \hat{X}(T_J)$. Then $\phi_{J,x}^* \mathcal{L}_{\xi} = \mathcal{L}_{J,x,\xi}$ is a (tame) local system on $G_{J,x}$ and $(\phi'_{J,x})^* \mathcal{L}_{\delta(\xi)} = \mathcal{L}'_{J,x,\xi}$ is a (tame) local system on $G'_{J,x}$. We denote by $A_{J,x,\xi}$ the perverse extension of $\mathcal{L}_{J,x,\xi}$ to $G/Z^0(L_J)$ and $A'_{J,x,\xi}$ the perverse extension of $\mathcal{L}'_{J,x,\xi}$ to $G/Z^0(L_{\delta(J)})$. We simply write $\mathcal{L}_{I,x,\xi}$ as $\mathcal{L}_{x,\xi}$ and $A_{I,x,\xi}$ as $A_{x,\xi}$.

By [MS1, 4.1.2], $\mathcal{L}_{J,x,\xi}$ and $A_{J,x,\xi}$ have weight $x\xi$ for the left B -action and $-\xi$ for the right B_J -action. Similarly, $\mathcal{L}'_{J,x,\xi}$ and $A'_{J,x,\xi}$ have weight $x\delta(\xi)$ for the left B -action and $-\delta(\xi)$ for the right $B_{\delta(J)}$ -action.

3.4. The group B acts on $G \times G/Z^0(L_J)$ by $b(g, h) = (gb^{-1}, bh)$. A quotient $G \times^B G/Z^0(L_J)$ exists and the product map $G \times G/Z^0(L_J) \rightarrow G/Z^0(L_J)$ defined by $(g, h) \mapsto gh$ induces a proper morphism

$$m_J : G \times^B G/Z^0(L_J) \rightarrow G/Z^0(L_J).$$

It is easy to see that $(m_J)_!(A_{w,\xi} \odot A_{J,1,\xi}) = A_{J,w,\xi}$.

For $\xi \in \hat{X}(T_J)$ and $w, x \in W$, set

$$\mathcal{I}(\xi, J, w, x) = \{w' \mid w'\xi = wx\xi \text{ and } G_{J,w'} \subset \overline{m_J(G_{I,w} \times G_{J,x})}\},$$

where $\overline{m_J(G_{I,w} \times G_{J,x})}$ is the closure of $m_J(G_{I,w} \times G_{J,x})$ in $G/Z^0(L_J)$. Then by [MS1, 4.2],

$$(m_J)_!(A_{w,x\xi} \odot A_{J,x,\xi}) \in < A_{J,w',\xi} >_{w' \in \mathcal{I}(\xi, J, w, x)}.$$

Similarly, set

$$\mathcal{I}'(\xi, J, w, x) = \{w' \mid w'\delta(\xi) = wx\delta(\xi) \text{ and } G'_{J,w'} \subset \overline{m_J(G_{I,w} \times G'_{J,x})}\}.$$

Then

$$(m_J)!(A_{w,x\xi} \odot A'_{J,x,\xi}) \in \langle A'_{J,w',\xi} \rangle_{w' \in \mathcal{I}'(\xi, J, w, x)}.$$

3.5. The group B_J acts on $G/Z^0(L_{\delta(J)}) \times G/Z^0(L_J)$ by $b(g, g') = (gg_0b^{-1}g_0^{-1}, g'b^{-1})$. The quotient $G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J)$ exists and the map $G \times G \rightarrow \overline{G^1}$ defined by $(g, g') \mapsto (g, g') \cdot h_J$ induces a morphism

$$p_J : G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J) \rightarrow \overline{G^1}.$$

For $w \in W^{\delta(J)}$ and $v \in W$, we have that

$$\begin{aligned} (B\dot{w}U_{P_{\delta(J)}^-}(B \cap L_{\delta(J)}), B\dot{v}B) \cdot h_J &= (B\dot{w}(B \cap L_{\delta(J)}), B\dot{v}(B \cap L_J)) \cdot h_J \\ &= (B\dot{w}(B \cap L_{\delta(J)}), B\dot{v}) \cdot h_J \\ &= (B\dot{w}, B\dot{v}) \cdot h_J = [J, w, v]. \end{aligned}$$

We can see that $p_J|_{G'_{J,w} \times^{B_J} G_{J,v}}$ factors through an affine space bundle map $p_{J,w,v} : G'_{J,w} \times^{B_J} G_{J,v} \rightarrow [J, w, v]$. For $\xi \in \hat{X}(T_J)$, $\mathcal{L}'_{J,w,\xi} \odot \mathcal{L}_{J,v,-\xi}$ is a local system on $G'_{J,w} \times^{B_J} G_{J,v}$ and

$$(p_J|_{G'_{J,w} \times^{B_J} G_{J,v}})!(\mathcal{L}'_{J,w,\xi} \odot \mathcal{L}_{J,v,-\xi}) = (i_{J,w,v})!\mathcal{L}_{\xi,J,w,v}.$$

For $w \in W^{\delta(J)}$ and $u \in W_J$, $p_J(G'_{J,w\delta(u)} \times^{B_J} G_{J,1}) = [J, w, u^{-1}]$ and $p_J|_{G'_{J,w\delta(u)} \times^{B_J} G_{J,1}}$ factors through an affine space bundle map $p_{J,w\delta(u),1} : G'_{J,w\delta(u)} \times^{B_J} G_{J,1} \rightarrow [J, w, u^{-1}]$. For $\xi \in \hat{X}(T_J)$, we have that

$$\begin{aligned} (i_{J,w,u^{-1}})!\mathcal{L}_{u\xi,J,w,u^{-1}} &= (p_J|_{G'_{J,w\delta(u)} \times^{B_J} G_{J,1}})!(\mathcal{L}'_{J,w\delta(u),\xi} \odot \mathcal{L}_{J,1,-\xi}) \\ &= (p_J|_{G'_{J,w} \times^{B_J} G_{J,u^{-1}}})!(\mathcal{L}'_{J,w,u\xi} \odot \mathcal{L}_{J,u^{-1},-u\xi}). \end{aligned}$$

3.6. The group B acts on $G \times \overline{G^1}$ by $b(g, z) = (gb^{-1}, (b, b)z)$ for $b \in B, g \in G$ and $z \in \overline{G^1}$. The quotient $G \times_B \overline{G^1}$ exists. The map $G \times \overline{G^1} \rightarrow \overline{G^1}$ defined by $(g, z) \mapsto (g, g) \cdot z$ induces a morphism $\rho : G \times_B \overline{G^1} \rightarrow \overline{G^1}$. The map ρ is the unique extension of the map γ defined in 2.3.

For $J \subset I$ and $w, v \in W$, we denote by $\rho_{J,w,v}$ the restriction of ρ to $G \times_B [J, w, v]$.

Let $\xi \in \hat{X}(T_J) \subset \hat{X}$ with $w\delta(\xi) = v\xi$. Then $\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi,J,w,v}$ is a local system on $G \times_B [J, w, v]$ and $\bar{\mathbb{Q}}_l[\dim(G)] \odot A_{\xi,J,w,v}$ is a simple perverse sheaf on $G \times_B \overline{G^1}$. Set $L_{\xi,J,w,v} = (\rho|_{G \times_B [J,w,v]})!(\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi,J,w,v})$ and $C_{\xi,J,w,v} = \rho_!(\bar{\mathbb{Q}}_l \odot A_{\xi,J,w,v})$. Then $L_{\xi,J,w,v}, C_{\xi,J,w,v} \in \mathcal{D}(\overline{G^1})$. Moreover, by the decomposition theorem in [BBD], $C_{\xi,J,w,v}$ is semisimple.

The following result is an easy consequence of 3.2.

Corollary 3.7. *We keep the notation of 3.2. Then*

$$\langle C_{\eta,J',w',v'} \rangle_{(J',w',v',\eta) \in \mathcal{I}_{J,w,v,\xi}} = \langle L_{\eta,J',w',v'} \rangle_{(J',w',v',\eta) \in \mathcal{I}_{J,w,v,\xi}}.$$

3.8. Similarly, B acts on $G \times (G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J))$ by $b(g, z) = (gb^{-1}, b \cdot z)$ for $b \in B$, $g \in G$ and $z \in G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J)$, where B acts diagonally on $G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J)$ on both factors on the left. Denote by $G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J))$ the quotient space.

The morphism $G \times G \times G \rightarrow \overline{G^1}$ defined by $(g, g_1, g_2) \mapsto (gg_1, \overline{gg_2}) \cdot h_J$ induces a morphism $\pi_J : G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J)) \rightarrow \overline{G^1}$. We simply write $\pi_J|_{G \times_B (G'_{J,w} \times^{B_J} G_{J,v})}$ as $\pi_{J,w,v}$. For $\xi \in \hat{X}(T_J)$ with $w\delta(\xi) = v\xi$, $\bar{\mathbb{Q}}_l \odot (\mathcal{L}'_{J,w,\xi} \odot \mathcal{L}_{J,v,-\xi})$ is a local system on $G \times_B (G'_{J,w} \times^{B_J} G_{J,v})$ and

$$(\pi_{J,w,v})_! (\bar{\mathbb{Q}}_l \odot (\mathcal{L}'_{J,w,\xi} \odot \mathcal{L}_{J,v,-\xi})) = L_{\xi,J,w,v}.$$

Lemma 3.9. *Let $J \subset I$, $w, v \in W$ and $\xi \in \hat{X}(T_J)$ with $w\delta(\xi) = v\xi$. Then*

$$(\pi_J)_! (\bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot A_{J,v,-\xi})) \in < (\pi_J)_! (\bar{\mathbb{Q}}_l \odot (A'_{J,w',\xi} \odot A_{J,1,-\xi})) >_{w' \in \mathcal{I}'(\xi, J, v^{-1}, w)}.$$

Proof. The group B_J acts on $G \times G/Z^0(L_J)$ on the second factor on the right. This action induces a B_J action on $G \times^B G/Z^0(L_J)$. Define the B_J action on $G/Z^0(L_{\delta(J)}) \times (G \times^B G/Z^0(L_J))$ by $b(g, z) = (gg_0b^{-1}g_0^{-1}, bz)$ for $g \in G/Z^0(L_{\delta(J)})$ and $z \in G \times^B G/Z^0(L_J)$. The quotient $G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J))$ exists and the morphism $(id, m_J) : G/Z^0(L_{\delta(J)}) \times (G \times^B G/Z^0(L_J)) \rightarrow G/Z^0(L_{\delta(J)}) \times G/Z^0(L_J)$ induces a morphism

$$G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J)) \rightarrow G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J).$$

The group B action on $G \times G/Z^0(L_J)$ by acting on the first factor on the left induces a B action on $G \times^B G/Z^0(L_J)$. Then we obtain the diagonal B -action on $G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J))$. We may also define the B -action on $G \times (G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J)))$ in the same way as we did in 3.8. We write the quotient space as $G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J)))$. The morphism $(id, id, m_J) : G \times G/Z^0(L_{\delta(J)}) \times (G \times^B G/Z^0(L_J)) \rightarrow G \times G/Z^0(L_{\delta(J)}) \times G/Z^0(L_J)$ induces a morphism

$$m_{1,2,34} : G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J))) \rightarrow G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J)).$$

The morphism $G \times G/Z^0(L_{\delta(J)}) \times G \times G/Z^0(L_J) \rightarrow \overline{G^1}$ defined by $(g_1, g_2, g_3, g_4) \mapsto (g_1g_2, g_1g_3g_4) \cdot h_J$ induces a morphism

$$f_{1,2,34} : G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J))) \rightarrow \overline{G^1}$$

and we have that $f_{1,2,34} = \pi_J \circ m_{1,2,34}$. We also have that

$$(m_{1,2,34})_! (\bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot (A_{v,-\xi} \odot A_{J,1,-\xi}))) = \bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot A_{J,v,-\xi})[\cdot].$$

Hence

$$(\pi_J)_! (\bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot A_{J,v,-\xi})) = (f_{1,2,34})_! (\bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot (A_{v,-\xi} \odot A_{J,1,-\xi})))[\cdot].$$

Similarly, we may define $G \times_B ((G \times^B G/Z^0(L_{\delta(J)})) \times^{B_J} G/Z^0(L_J))$ and the morphism $(id, m_{\delta(J)}, id) : G \times (G \times^B G/Z^0(L_{\delta(J)})) \times G/Z^0(L_J) \rightarrow G \times G/Z^0(L_{\delta(J)}) \times G/Z^0(L_J)$ induces a morphism

$$m_{1,23,4} : G \times_B ((G \times^B G/Z^0(L_{\delta(J)})) \times^{B_J} G/Z^0(L_J)) \rightarrow G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} G/Z^0(L_J)).$$

The morphism $G \times G \times G/Z^0(L_{\delta(J)}) \times G/Z^0(L_J) \rightarrow \overline{G^1}$ defined by $(g_1, g_2, g_3, g_4) \mapsto (g_1 g_2 g_3, g_1 g_4) \cdot h_J$ induces a morphism

$$f_{1,23,4} : G \times_B ((G \times^B G/Z^0(L_{\delta(J)})) \times^{B_J} G/Z^0(L_J)) \rightarrow \overline{G^1}$$

and we have that $f_{1,23,4} = \pi_J \circ m_{1,23,4}$.

The isomorphism $G \times G/Z^0(L_{\delta(J)}) \times G \times G/Z^0(L_J) \rightarrow G \times G \times G/Z^0(L_{\delta(J)}) \times G/Z^0(L_J)$ defined by $(g_1, g_2, g_3, g_4) \mapsto (g_1 g_3, g_3^{-1}, g_2, g_4)$ induces an isomorphism

$$\iota : G \times_B (G/Z^0(L_{\delta(J)}) \times^{B_J} (G \times^B G/Z^0(L_J))) \xrightarrow{\simeq} G \times_B ((G \times^B G/Z^0(L_{\delta(J)})) \times^{B_J} G/Z^0(L_J)).$$

It is easy to see that $f_{1,2,34} = f_{1,23,4} \circ \iota$ and

$$\iota! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot (A_{v,-\xi} \odot A_{J,1,-\xi})) \right) = \left(\bar{\mathbb{Q}}_l \odot ((A_{v^{-1},v\xi} \odot A'_{J,w,\xi}) \odot A_{J,1,-\xi}) \right) [\cdot].$$

Therefore

$$\begin{aligned} (\pi_J)! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w,\xi} \odot A_{J,v,-\xi}) \right) &= (f_{1,23,4})! \left(\bar{\mathbb{Q}}_l \odot ((A_{v\xi,v^{-1}} \odot A'_{J,w,\xi}) \odot A_{J,1,-\xi}) \right) [\cdot] \\ &= (\pi_J)! (m_{1,23,4})! \left(\bar{\mathbb{Q}}_l \odot ((A_{v\xi,v^{-1}} \odot A'_{J,w,\xi}) \odot A_{J,1,-\xi}) \right) [\cdot] \\ &\in < (\pi_J)! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w',\xi} \odot A_{J,1,-\xi}) \right) >_{w' \in \mathcal{I}'(\xi, J, v^{-1}, w)}. \end{aligned}$$

Corollary 3.10. *Let $J \subset I$, $w, v \in W$ and $\xi \in \hat{X}(T_J)$ with $w\delta(\xi) = v\xi$. Then*

$$L_{\xi, J, w, v} \in < L_{\xi, J, w', 1} >_{w' \in \mathcal{I}'(\xi, J, v^{-1}, w)}.$$

Proof. We have that

$$(\pi_{J,w,v})! \left(\bar{\mathbb{Q}}_l \odot (\mathcal{L}'_{J,w,\xi} \odot \mathcal{L}_{J,v,-\xi}) \right) \in < (\pi_J)! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w_1,\xi} \odot A_{J,v_1,-\xi}) \right) >_{w_1, v_1},$$

where w_1 runs over the elements in W such that G'_{J,w_1} is contained in the closure of $G'_{J,w}$ and v_1 runs over the elements in W such that G_{J,v_1} is contained in the closure of $G_{J,v}$.

By lemma 3.9, we have that

$$(\pi_J)! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w_1,\xi} \odot A_{J,v_1,-\xi}) \right) \in < (\pi_J)! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w_2,\xi} \odot A_{J,1,-\xi}) \right) >_{w_2},$$

where w_2 runs over the elements in W such that $w_2\delta(\xi) = \xi$ and G'_{J,w_2} is contained in the closure of $m_{\delta(J)}(G_{I,v_1^{-1}} \times G'_{J,w_1})$. Therefore,

$$\begin{aligned} (\pi_{J,w,v})! \left(\bar{\mathbb{Q}}_l \odot (\mathcal{L}'_{J,w,\xi} \odot \mathcal{L}_{J,v,-\xi}) \right) &\in < (\pi_J)! \left(\bar{\mathbb{Q}}_l \odot (A'_{J,w',\xi} \odot A_{J,1,-\xi}) \right) >_{w'} \\ &= < (\pi_{J,w',1})! \left(\bar{\mathbb{Q}}_l \odot (\mathcal{L}'_{J,w',\xi} \odot \mathcal{L}_{J,1,-\xi}) \right) >_{w'}, \end{aligned}$$

where w' runs over the elements in W such that $w'\delta(\xi) = \xi$ and $G'_{J,w'}$ is contained in the closure of $m_{\delta(J)}(G_{I,v^{-1}} \times G'_{J,w})$. $\square$

Lemma 3.11. *Let $J \subset I$ and $\mathcal{O}$ be a W_J -orbit on $\hat{X}(T_J)$. Set*

$$\begin{aligned}\mathcal{I}_{J,\mathcal{O}} &= \{(\xi, w, v) \mid \xi \in \mathcal{O}, w, v \in W \text{ with } w\delta(\xi) = v\xi\}, \\ \mathcal{I}'_{J,\mathcal{O}} &= \{(\xi, vw, 1) \in \mathcal{I}_{J,\mathcal{O}} \mid w \in W^{\delta(J)}, v \in W_{I(J,w,\delta)}\}.\end{aligned}$$

Then

$$< L_{\xi,J,w,v} >_{(\xi,w,v) \in \mathcal{I}_{J,\mathcal{O}}} = < L_{\xi,J,vw,1} >_{(\xi,vw,1) \in \mathcal{I}'_{J,\mathcal{O}}}.$$

Proof. Set $J' = \{i \in I \mid \alpha_i = -w_0\alpha_{\delta(j)} \text{ for some } j \in J\}$ and $y = w_0w_0^{\delta(J)}$. Then $y\delta(J) = J'$.

Let $w \in W^{\delta(J)}$ and $u \in W_{\delta(J)}$. Then we have that $wy^{-1} \in W^{J'}$ and $l(wuy^{-1}) = l(wy^{-1}yuy^{-1}) = l(wy^{-1}) + l(yuy^{-1}) = l(wy^{-1}) + l(u)$. Thus

$$\begin{aligned}\dim(G'_{J,wu}) &= \dim(G_{J,wuy^{-1}}) = l(wuy^{-1}) + \dim(B/Z^0(L_J)) \\ &= l(wy^{-1}) + l(u) + \dim(B/Z^0(L_J)).\end{aligned}$$

For $a, b \in W$, we have that $\dim(m_{\delta(J)}(G_{I,a} \times G'_{J,b})) \leq l(a) + l(by^{-1}) + \dim(B/Z^0(L_J))$ and if the equality holds, then $m_{\delta(J)}(G_{I,a} \times G'_{J,b}) = G'_{J,ab}$.

Now by 3.10, $< L_{\xi,J,w,v} >_{(\xi,w,v) \in \mathcal{I}_{J,\mathcal{O}}} = < L_{\xi,J,w,1} >_{(\xi,w,1) \in \mathcal{I}_{J,\mathcal{O}}}$. Moreover, for $\xi \in \mathcal{O}$, $w \in W^{\delta(J)}$ and $u \in W_J$ with $w\delta(u)\delta(\xi) = \xi$, we have that $L_{\xi,J,w\delta(u),1} = L_{u\xi,J,w,u^{-1}} \leq < L_{u\xi,J,w',1} >_{w' \in \mathcal{I}'(u\xi,J,u,w)}$. By induction, it suffices to prove the following statement:

Let $(w_i, u_i, \xi_i)_{i \geq 1}$ be a sequence with $w_i \in W^{\delta(J)}$, $u_i \in W_J$, $\xi_i \in \hat{X}(T_J)$. If $w_1\delta(u_1)\delta(\xi_1) = \xi_1$ and for each $i \geq 1$, $\xi_{i+1} = u_i\xi_i$ and $w_{i+1}\delta(u_{i+1}) \in \mathcal{I}'(\xi_{i+1}, J, u_i, w_i)$, then for $n \gg 0$, $u_n \in W_{I(J,w_n,\delta)}$.

For each $i \geq 1$, we have that

$$\begin{aligned}l(w_{i+1}y^{-1}) + l(u_{i+1}) &= \dim(G'_{J,w_{i+1}\delta(u_{i+1})}) - \dim(B/Z^0(L_J)) \\ &\leq \dim(m_{\delta(J)}(G_{I,u_i} \times G'_{J,w_i})) - \dim(B/Z^0(L_J)) \\ &\leq l(w_iy^{-1}) + l(u_i).\end{aligned}$$

Moreover, if the equalities hold, then $G'_{J,w_{i+1}\delta(u_{i+1})} = G'_{J,u_iw_i}$ and $w_{i+1}\delta(u_{i+1}) = u_iw_i$.

Thus for $k \gg 0$, we have $l(w_ky^{-1}) + l(u_k) = l(w_{k+1}y^{-1}) + l(u_{k+1}) = \dots$ and $w_{i+1}\delta(u_{i+1}) = u_iw_i$ for all $i \geq k$. By [H2, 3.10], $w_{i+1} \geq w_i$ for $i \geq k$. Therefore, for $m \gg k$, we have that $w_m = w_{m+1} = \dots = w_\infty$ and $l(u_m) = l(u_{m+1}) = \dots$. Thus, $w_\infty^{-1}u_iw_\infty = \delta(u_{i+1}) \in W_{\delta(J)}$ and $l(w_\infty^{-1}u_iw_\infty) = l(u_{i+1}) = l(u_i)$ for $i \geq m$. So $w_\infty^{-1}\text{supp}(u_i) = \delta(\text{supp}(u_{i+1}))$ for all $i \geq m$. In other words, for $i \geq m$,

$$\cup_{l \geq i+1} \text{supp}(u_l) \subset \cup_{l \geq i} \text{supp}(u_l) = w_\infty \delta(\cup_{l \geq i+1} \text{supp}(u_l)).$$

Thus for $n \gg m$, $\cup_{i \geq n} \text{supp}(u_i) = w_\infty \delta(\cup_{i \geq n} \text{supp}(u_i))$. So $\text{supp}(u_n) \subset \cup_{i \geq l} \text{supp}(u_i) \subset I(J, w_\infty, \delta)$ and $u_n \in W_{I(J,w_n,\delta)}$. The lemma is proved. $\square$

Combining 3.7 with 3.11, we obtain the key lemma.

Lemma 3.12. *Let $\mathcal{O}$ be a W -orbit on $\hat{X}$. Set*

$$\mathcal{I}_{\mathcal{O}} = \{(\xi, J, w, v) \mid J \subset I, \xi \in \mathcal{O} \cap \hat{X}(T_J), w, v \in W \text{ with } w\delta(\xi) = v\xi\},$$

$$\mathcal{I}'_{\mathcal{O}} = \{(\xi, J, vw, 1) \in \mathcal{I}_{\mathcal{O}} \mid w \in W^{\delta(J)}, v \in W_{I(J, w, \delta)}\}.$$

Then

$$< C_{\xi, J, w, v} >_{(\xi, J, w, v) \in \mathcal{I}_{\mathcal{O}}} = < L_{\xi, J, w, v} >_{(\xi, J, w, v) \in \mathcal{I}_{\mathcal{O}}} = < L_{\xi, J, vw, 1} >_{(\xi, J, vw, 1) \in \mathcal{I}'_{\mathcal{O}}}.$$

4. THE CHARACTER SHEAVES ON $\overline{G^1}$

By 2.1 and 3.12, we have the following result.

Proposition 4.1. *Let $\mathcal{O}$ be a W -orbit on $\hat{X}$ and A be a simple perverse sheaf on $\overline{G^1}$. The following conditions on A are equivalent:*

- (i) $A \dashv C_{\xi, J, w, v}$ for some $J \subset I$, $w, v \in W$ and $\xi \in \mathcal{O} \cap \hat{X}(T_J)$ with $w\delta(\xi) = v\xi$.
- (ii) $A \dashv L_{\xi, J, w, v}$ for some $J \subset I$, $w, v \in W$ and $\xi \in \mathcal{O} \cap \hat{X}(T_J)$ with $w\delta(\xi) = v\xi$.
- (iii) $A \dashv L_{\xi, J, vw, 1}$ for some $J \subset I$, $w \in W^{\delta(J)}$, $v \in W_{I(J, w, \delta)}$ and $\xi \in \mathcal{O} \cap \hat{X}(T_J)$ with $vw\delta(\xi) = \xi$.

4.2. Let $\mathcal{C}_{\mathcal{O}}(\overline{G^1})$ be the set of (isomorphism classes) of simple perverse sheaves on $\overline{G^1}$ which satisfies the equivalent conditions 4.1(i)-(iii) with respect to $\mathcal{O}$. The simple perverse sheaves on $\overline{G^1}$ which belong to $\mathcal{C}_{\mathcal{O}}(\overline{G^1})$ for some W -orbit $\mathcal{O}$ are called character sheaves on the group compactification $\overline{G^1}$; they (or their isomorphism classes) form a set $\mathcal{C}(\overline{G^1})$.

4.3. We keep the notation of 1.4. Now $K = I(J, w, \delta)$. Set $L = L_K/Z^0(L_J)$ and $B_1 = (B \cap L_K)/Z^0(L_J)$. For $v \in W_K$, set $L_v = B_1 \dot{v} \dot{w} g_0 B_1$.

Consider the following diagram

$$L \dot{w} g_0 \xleftarrow{p_1} G \times L \dot{w} g_0 \xrightarrow{p_2} G \times_{L_K} L \dot{w} g_0 \xrightarrow{\pi_J^w} Z_J^w$$

where p_1 is the projection to the second factor, p_2 is the projection map and π_J^w is the map in 1.6.

For any character sheaf X on $L_K \dot{w} g_0 / Z^0(L_J)$, let $\tilde{X}$ be the simple perverse sheaf on Z_J^w such that $\tilde{X} = (\pi_J^w)_!(\bar{\mathbb{Q}}_l \odot X)[\cdot]$. (Then $(\pi_J^w)^* \tilde{X}$ is a shift of $\bar{\mathbb{Q}}_l \odot X$. By the commuting diagram in 1.5, the simple perverse sheaf $\tilde{X}$ on Z_J^w is the same as the simple perverse sheaf obtained in [L3, 11.12].)

For $\xi \in \hat{X}(T_J)$, let $\mathcal{C}_{\xi}(Z_J^w)$ be the (isomorphism classes) of simple perverse sheaves on Z_J^w consisting of all $\tilde{X}$ as above for $X \in \mathcal{C}_{\xi}(L \dot{w} g_0)$. The simple perverse sheaves on Z_J^w which belong to $\mathcal{C}_{\xi}(Z_J^w)$ for some

$\xi \in \hat{X}(T_J)$ are called character sheaves on the G -stable piece Z_J^w ; they (or their isomorphism classes) form a set $\mathcal{C}(Z_J^w)$.

Now we study $\mathcal{C}_\xi(Z_J^w)$. For $v \in W_K$ with $vw\delta(\xi) = \xi$, let $\mathcal{L}_{\xi,v,w\delta}^L$ be the tame local system on L_v with weight $vw\delta(\xi)$ for the left B_1 -action and $A_{\xi,v,w\delta}^L$ be the perverse extension of $\mathcal{L}_{\xi,v,w\delta}^L$ to L .

We identify $L \times_{B_1} L_v$ with $L_K \times_{B \cap L_K} L_v$. Let $\bar{\mathbb{Q}}_l^G$ be the trivial local system on G and $\bar{\mathbb{Q}}_l^{L_K}$ be the trivial local system on L_K . Define $\rho_v^L : L_K \times_{B \cap L_K} L_v \rightarrow L\dot{w}g_0$ by $\rho(l, l') = ll'l^{-1}$. Then for $X \in \mathcal{C}_\xi(L\dot{w}g_0)$, there exists $v \in W_K$ with $vw\delta(\xi) = \xi$ and $X \dashv (\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot A_{\xi,v,w\delta}^L)$. By the decomposition theorem of [BBD], $(\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot A_{\xi,v,w\delta}^L)$ is semisimple. We have that $(\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot A_{\xi,v,w\delta}^L) = X[m] \oplus B$ for some $B \in \mathcal{D}(L)$ and $m \in \mathbb{Z}$. Therefore, $(\pi_J^w)_!\left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot A_{\xi,v,w\delta}^L))\right) = \tilde{X}[m'] \oplus C$ for some $C \in \mathcal{D}(Z_J^w)$ and $m' \in \mathbb{Z}$. In other words, a simple perverse sheaf A on Z_J^w is contained in $\mathcal{C}_\xi(Z_J^w)$ if and only if A is an irreducible constituent of $(\pi_J^w)_!\left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot A_{\xi,v,w\delta}^L))\right)$ for some $v \in W_K$ with $vw\delta(\xi) = \xi$.

Note that $\langle \mathcal{L}_{\xi,v,w\delta}^L \rangle_{v \in W_K, vw\delta(\xi)=\xi} = \langle A_{\xi,v,w\delta}^L \rangle_{v \in W_K, vw\delta(\xi)=\xi}$. Then a simple perverse sheaf A on Z_J^w is contained in $\mathcal{C}_\xi(Z_J^w)$ if and only if $A \dashv (\pi_J^w)_!\left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L))\right)$ for some $v \in W_K$ with $vw\delta(\xi) = \xi$.

4.4. We keep the notation of 4.3. Consider the commuting diagram

$$\begin{array}{ccccc} L_K \times_{B \cap L_K} L_v & \xleftarrow{p_1^L} & G \times (L_K \times_{B \cap L_K} L_v) & \xrightarrow{p_2^L} & G \times_{B \cap L_K} L_v \\ \downarrow \rho_v^L & & \downarrow (id, \rho_v^L) & & \downarrow \pi_v^L \\ L\dot{w}g_0 & \xleftarrow{p_1} & G \times L\dot{w}g_0 & \xrightarrow{p_2} & G \times_{L_K} L\dot{w}g_0, \end{array}$$

where p_1^L and p_2^L are projection maps and

$$\pi_v^L : G \times_{B_1} L_v = G \times_L (L \times_{B_1} L_v) \rightarrow G \times_L L\dot{w}g_0$$

is the map induced from (id, ρ_v^L) . Then

$$\begin{aligned} p_2^* \left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L)) \right) &= p_1^*(\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L) \\ &= (id, \rho_v^L)_!(p_1^L)^*(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L) = (id, \rho_v^L)_!(p_2^L)^*(\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L) \\ &= p_2^*(\pi_v^L)_!(\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L). \end{aligned}$$

Therefore $\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L)) = (\pi_v^L)_!(\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L)$.

Define $\pi_{B_1} : B/Z^0(L_J) \rightarrow B_1$ by $\pi_{B_1}(ub) = b$ for $u \in U_{P_{I(J,w,\delta)}}$ and $b \in B_1$. Define the map $B\dot{v}g_0(B \cap L_J)/Z^0(L_J) \rightarrow L_v$ by $b_1\dot{v}g_0b_2 \mapsto \pi_{B_1}(b_1)\dot{v}g_0\pi_{B_1}(b_2)$ for $b_1 \in B/Z^0(L_J)$ and $b_2 \in (B \cap L_J)/Z^0(L_J)$. It

is easy to see that this map is defined and is an affine space bundle morphism. This map induces an affine space morphism

$$p_v^L : G \times_{B \cap L_K} B\dot{w}g_0(B \cap L_J)/Z^0(L_J) \rightarrow G \times_{B \cap L_K} L_v.$$

We also have $(p_v^L)_!(p_v^L)^*(\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L) = (\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L)[\cdot]$.

Set $\phi = \pi_J^w \circ \pi_v^L \circ p_v^L : G \times_{B_1} B\dot{w}g_0(B \cap L_J)/Z^0(L_J) \rightarrow Z_J^w$. Then

$$\begin{aligned} (\pi_J^w)_! \left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L)) \right) &= (\pi_J^w)_! (\pi_v^L)_! (\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L) \\ &= \phi_! (p_v^L)^* (\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L)[\cdot]. \end{aligned}$$

Note that ϕ sends (g, g') to $(gg'g_0^{-1}, g) \cdot h_J$ for $g \in G$ and $g' \in B\dot{w}g_0(B \cap L_J)/Z^0(L_J)$. Now we identify $B\dot{w}g_0(B \cap L_J)/Z^0(L_J)$ with $B_1 \times_{B_1} B\dot{w}g_0(B \cap L_J)/Z^0(L_J)$. Then

$$\begin{aligned} \phi(B\dot{w}g_0(B \cap L_J)/Z^0(L_J)) &= (B\dot{w}(B \cap L_{\delta(J)}), 1) \cdot h_J \\ &= (B\dot{w}, B \cap L_J) \cdot h_J = [J, vw, 1]. \end{aligned}$$

By [H2, 1.12], $[J, vw, 1] \subset (P_K \dot{w}, P_K) \cdot h_J \subset Z_J^w$. Therefore

$$\rho(G \times_B [J, vw, 1]) \subset Z_J^w.$$

We denote by $\rho_J^{vw} : G \times_B [J, vw, 1] \rightarrow Z_J^w$ the restriction map of ρ .

Now the map $\phi : G \times_{B \cap L_K} G_{xw,\delta} \rightarrow Z_J^w$ factors through

$$G \times_{B \cap L_K} B\dot{w}g_0(B \cap L_J)/Z^0(L_J) \rightarrow G \times_{B \cap L_K} [J, vw, 1] \rightarrow G \times_B [J, vw, 1] \xrightarrow{\rho_J^{vw}} Z_J^w,$$

where the first two maps are affine space bundle maps. Hence

$$\begin{aligned} (\pi_J^w)_! \left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L)) \right) &= \phi_! (p_v^L)^* (\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,v,w\delta}^L)[\cdot] \\ &= (\rho_J^{vw})_! (\bar{\mathbb{Q}}_l^G \odot \mathcal{L}_{\xi,J,vw,1})[\cdot] = L_{\xi,J,vw,1} |_{Z_J^w} [\cdot]. \end{aligned}$$

Since $\rho(G \times_B [J, vw, 1]) \subset Z_J^w$, we have $L_{\xi,J,vw,1} |_{Z_{J'}^{w'}} = 0$ if $J \neq J'$ or $w \neq w'$.

In summary, for $(\xi, J, w, v) \in \mathcal{I}'_{\mathcal{O}}$,

$$L_{\xi,J,vw,1} |_{Z_{J'}^{w'}} = \begin{cases} (\pi_J^w)_! \left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_!(\bar{\mathbb{Q}}_l^{L_K} \odot \mathcal{L}_{\xi,v,w\delta}^L)) \right) [\cdot], & \text{if } J = J' \text{ and } w = w'; \\ 0, & \text{otherwise.} \end{cases}$$

Now we will show that the character sheaves on $\overline{G^1}$ have the following “nice” property.

Corollary 4.5. *Let $J \subset I$, $w \in W^{\delta(J)}$ and $\mathcal{O}$ be a W -orbit on $\hat{X}$. Let $A \in \mathcal{C}_{\mathcal{O}}(\overline{G^1})$ and K be a simple perverse sheaf on Z_J^w . If $K \dashv A |_{Z_J^w}$, then $K \in \mathcal{C}_{\xi}(Z_J^w)$ for some $\xi \in \mathcal{O} \cap \hat{X}(T_J)$.*

Proof. We keep the notation of 3.12.

There exists $(\xi, J', w', v) \in \mathcal{I}_{\mathcal{O}}$, such that $A \dashv C_{\xi,J',w',v}$. Since $C_{\xi,J',w',v}$ is semisimple, we have that $C_{\xi,J',w',v} = A[m] \oplus B$ for some $B \in \mathcal{D}(\overline{G^1})$ and $m \in \mathbb{Z}$. Then $C_{\xi,J',w',v} |_{Z_J^w} = A[m] |_{Z_J^w} \oplus B |_{Z_J^w}$. Thus $K \dashv C_{\xi,J',w',v} |_{Z_J^w}$.

By 2.1 and 3.12, $\langle C_{\xi, J', w', v} | Z_J^w \rangle_{(\xi, J', w', v) \in \mathcal{I}_O} = \langle L_{\xi, J', vw, 1} | Z_J^w \rangle_{(\xi, J', w', v) \in \mathcal{I}'_O}$. So $K \dashv (\pi_J^w)_! \left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_! (\bar{\mathbb{Q}}_l^{L_{I(J, w, \delta)}} \odot \mathcal{L}_{\xi, v, w\delta}^L)) \right)$ for some $v \in W_{I(J, w, \delta)}$ and $\xi \in \mathcal{O} \cap \hat{X}(T_J)$ with $vw\delta(\xi) = \xi$. By 4.3, $K \in \mathcal{C}_\xi(Z_J^w)$. The lemma is proved. $\square$

Corollary 4.6. *Let A be a simple perverse sheaf on $\overline{G^1}$. Then $A \in \mathcal{C}(\overline{G^1})$ if and only if there exists $J \subset I$, $w \in W^{\delta(J)}$ and $X \in \mathcal{C}(Z_J^w)$, such that A is the perverse extension of X .*

Remark. Therefore, our definition of character sheaves on $\overline{G^1}$ coincides with Lusztig's definition in [L3, 12.3].

Proof. We follow the proof of [L3, 11.15 & 11.18].

Since $\overline{G^1} = \bigsqcup_{J \subset I} \bigsqcup_{w \in W^{\delta(J)}} Z_J^w$, there exists $J \subset I$ and $w \in W^{\delta(J)}$, such that $\text{supp}(A) \cap Z_J^w$ is dense in $\text{supp}(A)$. Then $A|_{Z_J^w}$ is a simple perverse sheaf on Z_J^w and A is the perverse extension of $A|_{Z_J^w}$. By 4.5, $A|_{Z_J^w} \in \mathcal{C}(Z_J^w)$.

On the other hand, if $X \in \mathcal{C}(Z_J^w)$, then by 4.4, we may assume that $X \dashv (\rho_J^{vw})_! (\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi, J, vw, 1})$ for some $\xi \in \hat{X}(T_J)$ with $vw\delta(\xi) = \xi$. Then there exists a simple perverse sheaf A on $\overline{G^1}$ such that $\text{supp}(A)$ is the closure of $\text{supp}(X)$ in $\overline{G^1}$, $A|_{Z_J^w} = X$ and $A \dashv \rho_! (\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi, J, vw, 1})$. Then A is the perverse extension of X and $A \in \mathcal{C}(\overline{G^1})$. $\square$

Corollary 4.7. *Let $\mathcal{O}_1, \mathcal{O}_2$ be two distinct W -orbits on $\hat{X}$. Then for $(\xi_1, J_1, w_1, v_1) \in \mathcal{I}_{\mathcal{O}_1}$ and $(\xi_2, J_2, w_2, v_2) \in \mathcal{I}_{\mathcal{O}_2}$, we have that*

$$H_c(\overline{G^1}, C_{\xi_1, J_1, w_1, v_1} \otimes C_{-\xi_2, J_2, w_2, v_2}) = 0.$$

Proof. We keep the notation of 4.3. For any W_K -orbit $\mathcal{O}$ on $\hat{X}(T_J)$, set

$$\mathcal{I}_O^L = \{(v, \xi) \mid v \in W_K, \xi \in \mathcal{O} \text{ with } vw\delta(\xi) = \xi\}.$$

We have that $\langle \mathcal{L}_{\xi, v, w\delta}^L \rangle_{(v, \xi) \in \mathcal{I}_O^L} = \langle A_{\xi, v, w\delta}^L \rangle_{(v, \xi) \in \mathcal{I}_O^L}$. Now set $C_{\xi, v, w\delta}^L = (\pi_J^w)_! \left(\bar{\mathbb{Q}}_l^G \odot ((\rho_v^L)_! (\bar{\mathbb{Q}}_l^{L_K} \odot A_{\xi, v, w\delta}^L)) \right)$. Then $C_{\xi, v, w\delta}^L$ is semisimple and all the irreducible constituents are contained in $\mathcal{C}_\xi(Z_J^w)$. Moreover,

$$\langle L_{\xi, J, vw, 1} | Z_J^w \rangle_{(v, \xi) \in \mathcal{I}_O^L} = \langle C_{\xi, v, w\delta}^L \rangle_{(v, \xi) \in \mathcal{I}_O^L}.$$

Let $\mathcal{O}, \mathcal{O}'$ be two distinct W_K -orbits on $\hat{X}(T_J)$. Then by 2.6 and [MS1, 1.2.5], for $(v, \xi) \in \mathcal{I}_O^L$ and $(v', \xi') \in \mathcal{I}_{\mathcal{O}'}^L$, we have that

$$H_c(Z_J^w, C_{\xi, v, w\delta}^L \otimes C_{-\xi', v', w\delta}^L) = 0.$$

By 2.1, we also have that $H_c(Z_J^w, L_{\xi, J, vw, 1} | Z_J^w \otimes L_{-\xi', J, v'w, 1} | Z_J^w) = 0$.

By 4.4, for $(\xi_1, J_1, v_1 w_1, 1) \in \mathcal{I}_{\mathcal{O}_1}'$ and $(\xi_2, J_2, v_2 w_2, 1) \in \mathcal{I}_{\mathcal{O}_2}'$,

$$H_c(Z_J^w, L_{\xi_1, J_1, v_1 w_1, 1} | Z_J^w \otimes L_{-\xi_2, J_2, v_2 w_2, 1} | Z_J^w) = 0.$$

Since the above equality holds for all G -stable pieces Z_J^w of $\overline{G^1}$, we have that $H_c(\overline{G^1}, L_{\xi_1, J_1, v_1 w_1, 1} \otimes L_{-\xi_2, J_2, v_2 w_2, 1}) = 0$. Then by 2.1 and 3.12,

$$H_c(\overline{G^1}, C_{\xi_1, J_1, w_1, v_1} \otimes C_{-\xi_2, J_2, w_2, v_2}) = 0.$$

4.8. As in [L1, 11.12(c)], we have that $\mathcal{C}_{\mathcal{O}}(\overline{G^1}) \cap \mathcal{C}_{\mathcal{O}'}(\overline{G^1}) = \emptyset$ for distinct W -orbits $\mathcal{O}$ and $\mathcal{O}'$ on $\hat{X}$. In other words, there is a well defined map $\mathcal{C}(\overline{G^1}) \rightarrow W$ -orbit on $\hat{X}$ given by attaching $A \in \mathcal{C}(\overline{G^1})$ the W -orbit $\mathcal{O}$, where $A \in \mathcal{C}_{\mathcal{O}}(\overline{G^1})$.

5. THE PARABOLIC CHARACTER SHEAVES

In this section, G is a connected reductive algebraic group. We keep the notation in 1.3.

5.1. We first study the closures of the $B \times B$ -orbits in $\tilde{Z}_{J,y,\delta}$.

As in 3.5, we define the action of $B \cap L_J$ on $G/\dot{y}^{-1}U_{P_{J'}} \times G/U_{P_J}$ by $b(g, g') = (gg_0b^{-1}g_0^{-1}, g'b^{-1})$. Denote by $G/\dot{y}^{-1}U_{P_{J'}} \times^{B \cap L_J} G/U_{P_J}$ the quotient space. The morphism $G \times G \rightarrow \tilde{Z}_{J,y,\delta}$ defined by $(g_1, g_2) \mapsto (g_1, g_2) \cdot \tilde{h}_{J,y,\delta}$ induces a proper morphism

$$p_{J,y,\delta} : G/\dot{y}^{-1}U_{P_{J'}} \times^{B \cap L_J} G/U_{P_J} \rightarrow \tilde{Z}_{J,y,\delta}.$$

By 3.5, for $w \in W^{\delta(J)}$ and $v \in W$, we have that

$$p_{J,y,\delta}((B\dot{w}\dot{y}^{-1}B\dot{y})/\dot{y}^{-1}U_{P_{J'}} \times^{B \cap L_J} (B\dot{v}B)/U_{P_J}) = \widetilde{[J, w, v]_{y,\delta}}.$$

Moreover, for $w \in W^{\delta(J)}$, $u \in W_J$ and $v \in W$. we have that

$$\begin{aligned} (B\dot{w}\delta(u)\dot{y}^{-1}B\dot{y}, B\dot{v}B) \cdot \tilde{h}_{J,y,\delta} &= (B\dot{w}\delta(u)(B \cap L_{\delta(J)}), B\dot{v}B) \cdot \tilde{h}_{J,y,\delta} \\ &= (B\dot{w}, B\dot{v}B\dot{u}^{-1}) \cdot \tilde{h}_{J,y,\delta} \\ &\subset \bigcup_{v_1 \leq v, u_1 \leq u} (B\dot{w}, B\dot{v}_1\dot{u}_1^{-1}B) \cdot \tilde{h}_{J,y,\delta} \\ &= \bigcup_{v_1 \leq v, u_1 \leq u} \widetilde{[J, w, v_1 u_1^{-1}]_{y,\delta}}. \end{aligned}$$

Proposition 5.2. *Let $w \in W^{\delta(J)}$ and $v \in W$. Denote by $\widetilde{[J, w, v]_{y,\delta}}$ the closure of $[J, w, v]_{y,\delta}$ in $\tilde{Z}_{J,y,\delta}$. Then*

$$\overline{[J, w, v]_{y,\delta}} = \bigsqcup_{w_1 \in W^{\delta(J)}, u_1 \in W_J, w_1 \delta(u_1) y^{-1} \leq w y^{-1}} \bigsqcup_{v' \leq v} [J, w_1, v' u_1^{-1}]_{y,\delta}.$$

Proof. Note that $\bigsqcup_{w' \in W, w' y^{-1} \leq w y^{-1}} B\dot{w}'\dot{y}^{-1}B\dot{y}$ and $\bigsqcup_{v' \leq v} B\dot{v}'B$ are irreducible closed subvarieties of G and $p_{J,y,\delta}$ is a proper map. Thus we have that

$$\overline{[J, w, v]_{y,\delta}} = \bigcup_{w' y^{-1} \leq w y^{-1}, v' \leq v} p_{J,y,\delta}((B\dot{w}'\dot{y}^{-1}B\dot{y})/\dot{y}^{-1}U_{P_{J'}} \times^{B \cap L_J} (B\dot{v}'B)/U_{P_J}).$$

Let $w_1 \in W^{\delta(J)}$ and $u_1 \in W_J$. Then we have that $w_1 y^{-1} \in W^{J'}$ and $y\delta(u_1)y^{-1} \in W_{J'}$. Moreover, if $u_2 \leq u_1$, then $y\delta(u_2)y^{-1} \leq y\delta(u_1)y^{-1}$. Thus $w_1\delta(u_2)y^{-1} = w_1 y^{-1} y\delta(u_1)y^{-1} \leq w_1 y^{-1} y\delta(u_1)y^{-1} = w_1\delta(u_1)y^{-1}$. Therefore for $w_1\delta(u_1)y^{-1} \leq w y^{-1}$ and $v' \leq v$, we have that

$$\begin{aligned} & p_{J,y,\delta}((B\dot{w}_1\delta(\dot{u}_1)\dot{y}^{-1}B\dot{y})/\dot{y}^{-1}U_{P_{J'}} \times^{B \cap L_J} (B\dot{v}'B)/U_{P_J}) \\ & \subset \bigcup_{u_2 \in W_J, w_1\delta(u_2)y^{-1} \leq w_1\delta(u_1)y^{-1} \leq w y^{-1}} \bigcup_{v_1 \leq v' \leq v} \widetilde{[J, w_1, v_1 u_2^{-1}]_{y,\delta}}. \end{aligned}$$

Hence $\widetilde{[J, w, v]_{y,\delta}} \subset \bigsqcup_{w_1 \in W^{\delta(J)}, u_1 \in W_J, w_1\delta(u_1)y^{-1} \leq w y^{-1}} \bigsqcup_{v' \leq v} \widetilde{[J, w_1, v' u_1^{-1}]_{y,\delta}}$.

On the other hand, for $w_1 \in W^{\delta(J)}$, $u_1 \in W_J$ with $w_1\delta(u_1)y^{-1} \leq w y^{-1}$ and $v' \leq v$, we have that

$$\begin{aligned} \widetilde{[J, w_1, v' u_1^{-1}]_{y,\delta}} &= (B\dot{w}_1, B\dot{v}'\dot{u}_1^{-1}) \cdot \tilde{h}_{J,y,\delta} = (B\dot{w}_1\delta(\dot{u}_1), B\dot{v}') \cdot \tilde{h}_{J,y,\delta} \\ &\subset p_{J,y,\delta}((B\dot{w}_1\delta(\dot{u}_1)\dot{y}^{-1}B\dot{y})/\dot{y}^{-1}U_{P_{J'}} \times^{B \cap L_J} (B\dot{v}'B)/U_{P_J}) \\ &\subset \widetilde{[J, w, v]_{y,\delta}}. \end{aligned}$$

The lemma is proved. $\square$

5.3. As a consequence of the description of the closure of $B \times B$ -orbits, we will describe the closure of the G -stable pieces in $\tilde{Z}_{J,y,\delta}$ in 5.5. Although the result is not used in the proofs of properties of parabolic character sheaves we are going to discuss about, it serves as the motivation for them.

We have a bijection $\delta_y = y\delta : J \rightarrow J'$. Let $w, w' \in W$ with $l(w) = l(w')$. We say that w' can be obtained from w via a (J, δ_y) -cyclic shift if $w = s_{i_1} s_{i_2} \cdots s_{i_n}$ is a reduced expression and either (1) $i_1 \in J$ and $w' = s_{i_1} w s_{\delta_y(i_1)}$ or (2) $i_n \in J'$ and $w' = s_{\delta_y^{-1}(i_n)} w s_{i_n}$. We write $w \sim_{J,\delta_y} w'$ if there exists a finite sequence of elements $w = w_0, w_1, \dots, w_m = w'$ such that w_{k+1} can be obtained from w_k via a (J, δ_y) -cyclic shift.

By [H2, 3.9], we have the following result.

Lemma 5.4. *Let $w, w' \in W^{J'}$. Then the following conditions are equivalent:*

- (1) $w \geq u w' \delta_y(u)^{-1}$ for some $u \in W_J$.
- (2) $w \geq v w' \delta_y(u)^{-1}$ for some $v \leq u \in W_J$.
- (3) $w \geq x$ for some $x \sim_{J,\delta_y} w'$.

In this case, we say that $w \geq_{J,\delta_y} w'$.

Corollary 5.5. *Let $w \in W^{\delta(J)}$. Then the closure of $\tilde{Z}_{J,y,\delta}^w$ in $\tilde{Z}_{J,y,\delta}$ is $\bigsqcup_{w' \in W^{\delta(J)}, w y^{-1} \geq_{J,\delta_y} w' y^{-1}} \tilde{Z}_{J,y,\delta}^{w'}$.*

The case where $y = 1$ was proved in [H2, 4.6]. The general case can be treated in a similar way.

From now on, we study the parabolic character sheaves on $\tilde{Z}_{J,y,\delta}$.

5.6. Let $w, v \in W$. Let $p_{w,v} : B \times B \rightarrow [\widetilde{J, w, v}]_{y,\delta}$ be the morphism defined by $(b_1, b_2) \mapsto (b_1 \dot{w}, b_2 \dot{v}) \cdot \tilde{h}_{J,y,\delta}$ for $b_1, b_2 \in B$. By 1.3, $p_{wu,v\delta^{-1}(u)} = p_{w,v}$ for $u \in W_{\delta(J)}$.

The morphism $B \times B \rightarrow T$, $(t_1 u_1, t_2 u_2) \mapsto (\dot{w} g_0)^{-1} t_1 (\dot{w} g_0) \dot{v}^{-1} t_2^{-1} \dot{v}$ factors through a morphism $pr_{w,v} : [\widetilde{J, w, v}]_{y,\delta} \rightarrow T$.

For $\xi \in \hat{X}$, $pr_{w,v}^* \mathcal{L}_\xi = \mathcal{L}_{\xi,w,v}$ is a (tame) local system on $[\widetilde{J, w, v}]_{y,\delta}$. We denote by $A_{\xi,w,v}$ the perverse extension of $\mathcal{L}_{\xi,w,v}$ to $\tilde{Z}_{J,y,\delta}^w$.

5.7. We keep the notation of 1.4. Let $N_{\hat{G}}(L_K)$ be the normalizer of L_K in $\hat{G}$. Note that $N_{\hat{G}}(L_K)$ is a disconnected group with identity component L_K and $L_K \dot{w} g_0$ is a connected component of $N_{\hat{G}}(L_K)$. Consider the diagram

$$L_K \dot{w} g_0 \xleftarrow{p_1} G \times L_K \dot{w} g_0 \xrightarrow{p_2} G \times_{L_K} L_K \dot{w} g_0 \xrightarrow{\pi_{J,y,\delta}^w} \tilde{Z}_{J,y,\delta}^w$$

where p_1 is the projection to the second factor and p_2 is the projection map.

For any character sheaf X on $L_K \dot{w} g_0$, let $\tilde{X}$ be the simple perverse sheaf on $\tilde{Z}_{J,y,\delta}^w$ such that $\tilde{X} = (\tilde{\pi}_{J,y,\delta}^w)_!(\bar{\mathbb{Q}}_l \odot X)[\cdot]$. Let $\mathcal{C}(\tilde{Z}_{J,y,\delta}^w)$ be the (isomorphism classes) of simple perverse sheaves on $\tilde{Z}_{J,y,\delta}^w$ consisting of all $\tilde{X}$ as above. The elements in $\mathcal{C}(\tilde{Z}_{J,y,\delta}^w)$ are called the character sheaves on $\tilde{Z}_{J,y,\delta}^w$.

5.8. The group B acts on $G \times \tilde{Z}_{J,y,\delta}$ by $b(g, z) = (gb^{-1}, (b, b)z)$. The quotient $G \times_B \tilde{Z}_{J,y,\delta}$ exists. The map $G \times \tilde{Z}_{J,y,\delta} \rightarrow \tilde{Z}_{J,y,\delta}$ defined by $(g, z) \mapsto (g, g) \cdot z$ induces a morphism $\rho : G \times_B \tilde{Z}_{J,y,\delta} \rightarrow \tilde{Z}_{J,y,\delta}$.

By what we did in section 3 and section 4, we obtain the following result.

Proposition 5.9. *Let A be a simple perverse sheaf on $\tilde{Z}_{J,y,\delta}$. The following conditions are equivalent:*

- (i) $A \dashv \rho_!(\bar{\mathbb{Q}}_l \odot A_{\xi,w,v})$ for $w, v \in W$ and $\xi \in \hat{X}$ with $w\delta(\xi) = v\xi$.
- (ii) $A \dashv (\rho|_{G \times_B [\widetilde{J,w,v}]_{y,\delta}})_!(\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi,w,v})$ for some $w, v \in W$ and $\xi \in \hat{X}$ with $w\delta(\xi) = v\xi$.

- (iii) $A \dashv (\rho|_{G \times_B [\widetilde{J,vw,1}]_{y,\delta}})_!(\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi,vw,1})$ for some $w \in W^{\delta(J)}$, $v \in W_{I(J,w,\delta)}$ and $\xi \in \hat{X}$ with $vw\delta(\xi) = \xi$.

- (iv) There exists $w \in W^{\delta(J)}$ and $X \in \mathcal{C}(\tilde{Z}_{J,y,\delta}^w)$ such that A is the perverse extension of X to $\tilde{Z}_{J,y,\delta}$.

Moreover, if A satisfies the equivalent conditions, then for any $w \in W^{\delta(J)}$, any composition factor of ${}^p H^i(A|_{\tilde{Z}_{J,y,\delta}^w})$ is contained in $\mathcal{C}(\tilde{Z}_{J,y,\delta}^w)$.

Remark. The condition (iv) was one of the two equivalent definitions of parabolic character sheaves in [L3]. The fact the (i) $\Leftrightarrow$ (iv) was first

proved by Springer in [S2] in terms of (P, Q, σ) -character sheaves on G . The “moreover” part was first proved by Lusztig in [L3, 11.14]. Our approach gave a new proof of their results.

5.10. Let $\mathcal{C}(\tilde{Z}_{J,y,\delta})$ be the (isomorphism classes) of simple perverse sheaves on $\tilde{Z}_{J,y,\delta}$ which satisfy the equivalent conditions 5.9(i)-(iv). The elements of $\mathcal{C}(\tilde{Z}_{J,y,\delta})$ are called the parabolic character sheaves on $\tilde{Z}_{J,y,\delta}$.

For W_J -orbit $\mathcal{O}$ on $\hat{X}$, we denote by $\mathcal{C}_{\mathcal{O}}(\tilde{Z}_{J,y,\delta})$ the (isomorphism classes) of simple perverse sheaves A on $\tilde{Z}_{J,y,\delta}$ such that A satisfies the condition 5.9(iii) for some $\xi \in \mathcal{O}$. Then as in 4.8, we have that

$$\mathcal{C}_{\mathcal{O}}(\tilde{Z}_{J,y,\delta}) \cap \mathcal{C}_{\mathcal{O}'}(\tilde{Z}_{J,y,\delta}) = \emptyset$$

for distinct W_J -orbits $\mathcal{O}$ and $\mathcal{O}'$ on $\hat{X}$. In other words, there is a well defined map $\mathcal{C}(\tilde{Z}_{J,y,\delta}) \rightarrow W_J\text{-orbit on } \hat{X}$ given by attaching $A \in \mathcal{C}(\tilde{Z}_{J,y,\delta})$ the W -orbit $\mathcal{O}$, where $A \in \mathcal{C}_{\mathcal{O}}(\tilde{Z}_{J,y,\delta})$. This is a generalization of [L2, 5.3].

5.11. Now let us recall another definition of parabolic character sheaves in [L3].

Let $w \in W$. The Borel group B acts on $G_{I,w} \times G_{I,y}$ by $b(g, h) = (gb^{-1}, bh)$. The quotient $G_{I,w} \times^B G_{I,y}$ exists. The action of B on $G \times (G_{I,w} \times G_{I,y})$ defined by $b \cdot (g, g_1, g_2) = (gb^{-1}, bg_1, g_2g_0b^{-1}g_0^{-1})$ induces an action of B on $G \times (G_{I,x} \times^B G_{I,y})$. The quotient $G \times_B (G_{I,x} \times^B G_{I,y})$ exists.

With the notation of [L2, 4.2], set

$$Y_{(w,y)} = \{(B, B', g) \in \mathcal{B} \times \mathcal{B} \times G^1 \mid \text{pos}(B, B') = w, \text{pos}(B', {}^g B) = y\}.$$

The morphism $G \times (G_{I,w} \times G_{I,y}) \rightarrow Y_{(w,y)}$, $(g, g_1, g_2) \mapsto ({}^g B, {}^{gg_1} B, gg_1g_2g_0g^{-1})$ induces an isomorphism $G \times_B (G_{I,w} \times^B G_{I,y}) \cong Y_{(w,y)}$.

Let $\xi \in \hat{X}$ with $wy\delta(\xi) = \xi$. Then $\tilde{\mathcal{L}}_{\xi,w} = \bar{\mathbb{Q}}_l \odot (\mathcal{L}_{y\delta(\xi),w} \odot \mathcal{L}_{\delta(\xi),y})$ is a (tame) local system on $G \times_B (G_{I,w} \times^B G_{I,y})$. (The local system $\tilde{\mathcal{L}}$ on $Y_{(w,y)}$ defined in [L2, 4.2] is of the form $\tilde{\mathcal{L}}_{\xi,w}$ for some $\xi \in \hat{X}$ with $wy\delta(\xi) = \xi$ via the isomorphism $G \times_B (G_{I,w} \times^B G_{I,y}) \cong Y_{(w,y)}$.)

Let $\rho_{(w,y)} : G \times_B (G_x \times^B G_{y,\delta}) \rightarrow \tilde{Z}_{J,y,\delta}$ be the morphism defined by $(g, g_1, g_2) \mapsto ({}^g P_J, {}^{gg_1} P_{J'}, gg_1 U_{P_{J'}} g_2 g_0 U_P g^{-1})$. Let $\mathcal{C}_{J,y,\delta}$ be the set of (isomorphism classes of) simple perverse sheaves A on $\tilde{Z}_{J,y,\delta}$ such that $A \dashv (\rho_{(w,y)})^* \tilde{\mathcal{L}}_{\xi,w}$ for some $w \in W$ and $\xi \in \hat{X}$ with $wy\delta(\xi) = \xi$. This is the definition of parabolic character sheaves on $\tilde{Z}_{J,y,\delta}$ in [L3, 11.3].

Corollary 5.12. *We have that $\mathcal{C}(\tilde{Z}_{J,y,\delta}) = \mathcal{C}_{J,y,\delta}$.*

Remark. Thus we obtained a new proof of the fact that the two definitions of parabolic character sheaves in [L3] coincide, which was first proved by Lusztig in [L3, 11.15 & 11.18].

Proof. We identify $G_{I,w} \times^B G_{I,y}$ with $B \times_B (G_{I,w} \times^B G_{I,y})$. Then

$$\begin{aligned} \rho_{(w,y)}(G_{I,w} \times^B G_{I,y}) &= (P_J, {}^{B\dot{w}}P_{J'}, B\dot{w}B\dot{y}Bg_0) \\ &= (P_J, {}^{B\dot{w}\dot{y}\dot{y}^{-1}}P_{J'}, B\dot{w}\dot{y}U_{\dot{y}^{-1}P_{J'}}Bg_0) \\ &= [\widetilde{J, wy, 1}]_{y,\delta}. \end{aligned}$$

Let $\rho'_{(w,y)} : G_{I,w} \times^B G_{I,y} \rightarrow [\widetilde{J, wy, 1}]_{y,\delta}$ be the restriction of $\rho_{(w,y)}$. Then $\rho'_{(w,y)}$ is an affine space bundle map and $(\rho'_{(w,y)})_!(\mathcal{L}_{w,y\delta(\xi)} \odot \mathcal{L}_{y,\delta(\xi)}) = \mathcal{L}_{\xi,wy,1}$ for $\xi \in \hat{X}$. The morphism $(id, \rho'_{(w,y)}) : G \times (G_{I,w} \times^B G_{I,y}) \rightarrow G \times [\widetilde{J, wy, 1}]_{y,\delta}$ induces a morphism

$$\tilde{\rho}_{(w,y)} : G \times_B (G_{I,w} \times^B G_{I,y}) \rightarrow G \times_B [\widetilde{J, wy, 1}]_{y,\delta}.$$

We have that $\rho_{(w,y)} = \rho|_{G \times [\widetilde{J, wy, 1}]_{y,\delta}} \circ \tilde{\rho}_{(w,y)}$. Hence for $\xi \in \hat{X}$ with $wy\delta(\xi) = \xi$, we have that

$$(\rho_{(w,y)})_!\tilde{\mathcal{L}}_{\xi,w} = (\rho|_{G \times [\widetilde{J, wy, 1}]_{y,\delta}})_!(\tilde{\rho}_{(w,y)})_!\tilde{\mathcal{L}}_{\xi,w} = (\rho|_{G \times [\widetilde{J, wy, 1}]_{y,\delta}})_!(\bar{\mathbb{Q}}_l \odot \mathcal{L}_{\xi,wy,1})[\cdot].$$

Now let A be a simple perverse sheaf on $\tilde{Z}_{J,y,\delta}$. If A satisfies 5.9(iii), then $A \in \mathcal{C}_{J,y,\delta}$. If $A \in \mathcal{C}_{J,y,\delta}$, then A satisfies 5.9(ii). Therefore, $\mathcal{C}(\tilde{Z}_{J,y,\delta}) = \mathcal{C}_{J,y,\delta}$. $\square$

Note added. After completing this paper, I learned that the main result of this paper has also been obtained by T.A.Springer (unpublished).

REFERENCES

- [BBD] A. A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), Soc. Math. France, Paris, 1982, pp. 5–171.
- [H1] X. He, *Unipotent variety in the group compactification*, math.RT/0410199, accepted by Adv. in Math.
- [H2] X. He, *The G-stable pieces of the wonderful compactification*, math.RT/0412302, accepted by Trans. Amer. Math. Soc.
- [L1] G. Lusztig, *Character sheaves III*, Adv. in Math. **57** (1985), no. 3, 266–315.
- [L2] G. Lusztig, *Parabolic character sheaves, I*, Mosc. Math. J. **4** (2004), no. 1, 153–179.
- [L3] G. Lusztig, *Parabolic character sheaves, II*, Mosc. Math. J. **4** (2004), no. 4, 869–896.
- [MS1] J. G. M. Mars and T. A. Springer, *Character sheaves*, Astérisque no. 173–174 (1989), 9, 111–198.
- [MS2] J. G. M. Mars and T. A. Springer, *Hecke algebra representations related to spherical varieties*, Represent. Theory **2** (1998), 33–69.
- [S1] T. A. Springer, *Intersection cohomology of $B \times B$ -orbits closures in group compactifications*, J. Alg. **258** (2002), 71–111.
- [S2] T. A. Springer, *Remarks on parabolic character sheaves (preliminary version)*, unpublished.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY,
CAMBRIDGE, MA 02139, USA

E-mail address: `xuhua@mit.edu`